

## **Reduced melatonin synthesis in pregnant night workers: metabolic implications for offspring.**

Nehme P.A <sup>1</sup>, Amaral F <sup>2</sup>, Lowden A<sup>3</sup>, Skene D.J <sup>4</sup>, Cipolla-Neto J <sup>5</sup>, Moreno C.R.C<sup>1,3</sup>

1 School of Public Health, University of São Paulo

2 Department of Physiology, Federal University of São Paulo

3 Stress Research Institute, University of Stockholm

4 Faculty of Health and Medical Sciences, University of Surrey

5 Department of Physiology and Biophysics Neurobiology Lab, Institute of Biomedical Sciences, University of São Paulo

**Corresponding author at: crmoreno@usp.br**

### **Abstract**

Several novel animal studies have shown that intrauterine metabolic programming can be modified in the event of reduced melatonin synthesis during pregnancy, leading to glucose intolerance and insulin resistance in the offspring. It is therefore postulated that female night workers when pregnant may expose the offspring to unwanted health threats. This may be explained by the fact that melatonin is essential for regulating energy metabolism and can influence reproductive activity. Moreover, the circadian misalignment caused by shift work affects fertility and the fetus, increasing the risk of miscarriage, premature birth and low birth weight, phenomena observed in night workers. Thus, we hypothesize that light-induced melatonin suppression as a result of night work may alter intrauterine metabolic programming in pregnant women, potentially leading to metabolic disorders in their offspring.

Keywords: pregnancy, offspring, melatonin, metabolism, shift work.

## **Introduction**

Night workers experience a misalignment between their endogenous circadian timing system and behavior associated with work hours (mistimed sleep/wake; light/dark; feeding/fasting)[1–3]. In addition, light at night can acutely suppress melatonin production [4–6]. In this paper we focus on the possible risks threats to reproductive status in women undertaking night work and experiencing light-induced melatonin suppression [7]. Above all, we discuss metabolic implications for offspring due to maternal melatonin suppression.

Melatonin is an indoleamine synthesized in most animals and plants [8]. In mammals melatonin is produced primarily by the pineal gland. The light/dark cycle entrains the circadian oscillator in the hypothalamic suprachiasmatic nuclei (SCN), which in turn, entrains the pineal melatonin rhythm. Melatonin production occurs at night [9], and circulates from the pineal gland via the bloodstream and cerebrospinal fluid to other tissues of the body [10]. However, melatonin has several physiological actions besides signaling time of day or time of year information to the organism, such as its immunomodulatory [11], antioxidant [12] and cytostatic effects [13,14]. The light sensitivity of the system controls melatonin production, which together with the SCN, determines the duration of its synthesis [15]. Nevertheless, exposure to artificial light at night, as in the case of shift and night work, modifies melatonin production [16–18]. This occurs due to an acute effect of light at night on melatonin, light inhibiting its synthesis [19].

Although natural light differs to electric light, especially in terms of spectral composition and intensity, electric light is still a potent blocker of melatonin synthesis

[5,6]. Moreover, recent studies have demonstrated a lower threshold for light intensity to suppress melatonin than previously reported [20]. The exposure to light at night has been shown to reduce melatonin levels with consequent circadian misalignment in night workers as discussed below [16].

### **Night work and health**

Night work is common in contemporary society. The structure of modern society demands continuous production and/or services around the clock [21,22]. Across the world in industrialized societies it is estimated that approximately one in five workers work nights or irregular hours [23,24], where most are subjected to circadian misalignment where the behavioural cycles of sleep/wake and feeding/fasting are misaligned with the circadian timing system [25,26]. The circadian timing system is organized to facilitate behavioral and physiological adaptation (including sleep) to the 24-hour light-dark cycle [27]. In mammals, in addition to the “master” timing system located in the hypothalamic SCN, peripheral oscillators occur in the brain and throughout the body [28,29]. The SCN clock contributes to the regulation of body temperature, secretion of hormones melatonin and cortisol, sleep and wakefulness [30-32].

Studies investigating the effects of night work on workers’ health have revealed that abnormal working hours can pose a health risk by directly acting on metabolic processes, for instance affecting body mass regulation, gut efficiency and metabolic rate [33-36]. Detrimental effects have been observed in lipid and carbohydrate metabolism, as well as in insulin resistance [37,38]. The negative impacts of night work also include changes in sleep and wakefulness [23,39], and consequent alterations in meal timing [40–42] and physical activity [43,44]. Shift work has been strongly associated with

cardiovascular diseases [45], gastrointestinal [1] and metabolic disorders [46–48], and cancer [49]. Although there is less strong evidence linking shift work to mental health problems and reproduction-related problems, a few studies show its association with these factors such as depression [50], and occurrence of miscarriage [7], among other effects related to reproduction [7,51]. The social impact of changes in the pattern of family activities has also been reported [52].

Most of these adverse effects stem from misalignment of circadian timing and behavior (sleep/wake; feeding/fasting) defined by Erren et al. [25] as chronodisruption, a significant disturbance in circadian organization of the physiology, endocrinology, metabolism and behavior of living organisms, caused in part by exposure to light at night that leads to reduced melatonin synthesis [53].

### **Melatonin, shift work and reproductive activity**

Melatonin is essential for regulating energy metabolism [54] but also for reproductive activity, promoting changes in the gonadal pituitary axis and prolactin secretion [55] and consequently modulating reproductive physiology [56,57]. Increased melatonin levels are associated with delayed puberty, and hypothalamic amenorrhea whereas decreased melatonin is found in precocious puberty [58,59]. Thus, it is increasingly relevant to study the association of melatonin with improvements in ovulation rate, luteal function and increased embryo viability [60,61]. In addition, deficiencies in melatonin production during pregnancy may alter intrauterine metabolic programming, potentially leading to glucose intolerance and insulin resistance in offspring [61–64].

Besides the acute effects on melatonin production due to light at night, shift workers may exhibit circadian misalignment and sleep deprivation which may affect

fertility and the fetus, increasing the risk of miscarriage, premature birth and low birth weight [65,66]. Seron-Ferre et al. [67] analyzed the effects of circadian misalignment on the fetus and noted that this situation leads to alteration in the body temperature rhythm of newborns. Continuous exposure to light during pregnancy may increase the risk of spatial memory deficit in the adult life of the offspring in rats [68]. Most importantly, even under a light-dark cycle, the absence of melatonin in the mother during gestation impairs physical growth, neurodevelopment, and cognitive behavior in the offspring [69].

Notably, such factors (impaired physical growth, neurodevelopment, and cognitive behavior) are strong predictors of chronic diseases in adulthood. A relationship has been established between deleterious prenatal environment (e.g. malnutrition and/or fetal hypoxia) and the occurrence of diseases in adult life including hypertension, coronary heart disease, stroke, as well as metabolic and neurological disorders [70].

The mechanism by which melatonin plays a key role in the reproductive activity of numerous mammalian species [71] is via the transmission of photoperiodic information, which contributes to the seasonal modulation of reproductive activity [61]. Moreover, there is experimental evidence that maternal melatonin during pregnancy also provides a signal for the fetus [67,72]. Considering this signal follows a circadian rhythm, it suggests that a circadian pattern may be an important cue for fetal development as well to help the newborn to be entrained to the light-dark cycle after birth. Melatonin levels increase throughout the pregnancy, including in humans [73], which corroborates the idea of melatonin playing a role in fetal development.

Melatonin crosses the placenta freely and unaltered, entering fetal circulation and providing the fetus with photoperiodic signals [67,74,75]. Although circadian

rhythms are established during the fetal period [76], the rhythms are initially synchronized to the mother's circadian timing system. Moreover, human fetuses have also shown circadian rhythmicity in sleep/wake and other behaviors [72,77]. The diurnal rhythmicity, however, is interrupted when mothers are exposed to constant light [78], conditions associated with melatonin suppression.

### **Metabolic effects due to changes in melatonin rhythm**

As mentioned earlier, melatonin has an important role in optimizing energy balance and body weight regulation [79], crucial events for metabolic health[10,80]. Diseases such as type 2 diabetes, coronary artery disease, hypertension and obesity may be related to impaired intrauterine development. This phenomenon called "programming" or "hypothesis of fetal origin diseases" refers to permanent changes in the structure, physiology or metabolism of certain organs, caused by stimuli or diseases during the intrauterine development period [81,82].

Results of studies linking the external and internal environment during pregnancy to effects on body composition of the offspring highlight that this period is crucial in the genesis of obesity and possibly numerous other diseases [83,84]. In a study with rats, Diaz and Blasquez [85] concluded that the removal of the pineal had a significant impact on plasma glucose, insulin and glucagon secretions. These authors investigated three groups of animals, controls, pinealectomized rats and pinealectomized and melatonin-treated rats. Baseline glucose, insulin and glucagon levels were evaluated after administration of arginine and glucose. The animals were kept under controlled lighting conditions, with alternating cycles of 12 hours of light and dark. The results of this study indicated that the pinealectomized animals had the highest levels of glucose, the animals treated with melatonin had intermediary levels of

glucose and the lowest levels were found in the controls. In addition, pinealectomized rats showed changes in insulin levels, evidenced by changes in glycogenesis and glucose uptake, highlighting to inverse relationship between insulin and glucagon levels in pinealectomized animals. Finally, treatment with melatonin partially restored the levels of insulin and glucagon in pinelectomized animals emphasizing the importance of pineal to metabolism. However, the functional interplay between melatonin and insulin was demonstrated by Lima et al. [86]. In this study, the authors showed for the first time that the absence of melatonin lead to impaired glucose tolerance and a state of insulin resistance. Later on, the same group demonstrated that melatonin, acting through its MT1 and MT2 receptors, was able to phosphorylate the insulin receptor potentiating insulin action either in the central nervous system or in the periphery [87,88].

Given melatonin's key role in the homeostasis of carbohydrate metabolism, Van Cauter and others [89–91] elucidated the pathways by which pineal melatonin may act as a modulator of glucose metabolism, considering that there is no pattern in glucose tolerance over the day, either through food intake or another kind of glucose intake [90]. To reinforce such an hypothesis, studies with animal models have shown increased insulin levels during the day and reduced levels of insulin concomitant with elevated glucose levels at night, concurrently with melatonin elevation [92,93]. In addition, melatonin has a stimulatory effect on glucagon and pancreatic  $\alpha$  cells and, on the action of hepatic glucagon in the human liver [94,95]. Increased release of glucagon occurs in response to stimulation of human pancreatic cells which in turn promotes insulin secretion [96].

The action exerted by endogenous melatonin occurs through interaction with its membrane receptors, MT1 and MT2, on the cell. A reaction mediated by these receptors triggers the release of inhibitory G proteins, with consequent decrease in adenylyl-

cyclase activity and reduction of cAMP. Through the use of antagonists of these receptors (4-phenyl-2-propionamidotetralin and luzindole), the importance of melatonin in the regulation of leptin expression has been shown [97,98]. Animal models have shown that the influence of feeding behavior potentially promotes physical changes, such as affecting the fat-lean mass ratio, which could lead to obesity [99–101]. Appetite hormones such as leptin may be related to this process [102–104].

### **The hypothesis:**

The cited literature suggests that aspects of modern life, such as night work and shift work, may alter circadian rhythmicity and melatonin production [105,106] in key stages of human life, such as the reproductive stage, given the fact that pregnant women may also engage in night work. Although studies on fetal programming have been carried out using animal models, there is strong evidence of reduced melatonin production in night workers as well as negative health effects. We thus hypothesize that the light-induced suppression of melatonin in pregnant night workers may affect intrauterine metabolic programming, potentially leading to metabolic disorders in the offspring.

### **Conclusion:**

In summary, there is some evidence that shift work can affect fetal development. Melatonin in some models has been shown to affect reproductive function and is important in fetal development and metabolic programming. Shift workers are exposed to light at night, which in turn is known to suppress melatonin synthesis. Based on these assumptions, we have formulated our study hypothesis.



**Disclosure:**

The authors declare no conflict of interest.

**Funding:**

This work was supported by the Fundação de Amparo a Pesquisa do Estado de São Paulo – FAPESP (2014/50457-0 and 2016/11155-3); Coordenação de Aperfeiçoamento de Pessoal de Nivel Superior – CAPES– Finance Code 001; CAPES/Stint (021/14); and Conselho Nacional de Desenvolvimento Científico e Tecnológico -CNPq (142140/2015-5).

## REFERENCES

- [1] Knutsson A, Bøggild H. Gastrointestinal disorders among shift workers. *Scand J Work Environ Health* 2010;36:85–95. doi:10.5271/sjweh.2897.
- [2] McHill AW, Melanson EL, Higgins J, Connick E, Moehlman TM, Stothard ER, et al. Impact of circadian misalignment on energy metabolism during simulated nightshift work. *Proc Natl Acad Sci* 2014;111:17302–7. doi:10.1073/pnas.1412021111.
- [3] Potter GDM, Skene DJ, Arendt J, Cade JE, Grant PJ, Hardie LJ. Circadian rhythm and sleep disruption: Causes, metabolic consequences, and countermeasures. *Endocr Rev* 2016;37:584–608. doi:10.1210/er.2016-1083.
- [4] Zeitzer JM, Dijk DJ, Kronauer R, Brown E, Czeisler C. Sensitivity of the human circadian pacemaker to nocturnal light: melatonin phase resetting and suppression. *J Physiol* 2000;526 Pt 3:695–702. doi:10.1111/j.1469-7793.2000.00695.x.
- [5] Brainard GC, Hanifin JP, Greeson JM, Byrne B, Glickman G, Gerner E, et al. Action spectrum for melatonin regulation in humans: evidence for a novel circadian photoreceptor. *J Neurosci* 2001;21:6405–12. doi:10.1523/JNEUROSCI.2111-01.2001 [pii].
- [6] Thapan K, Arendt J, Skene DJ. An action spectrum for melatonin suppression: Evidence for a novel non-rod, non-cone photoreceptor system in humans. *J Physiol* 2001;535:261–7. doi:10.1111/j.1469-7793.2001.t01-1-00261.x.
- [7] Palmer KT, Bonzini M, Harris EC, Linaker C, Bonde JP. Europe PMC Funders Group Work activities and risk of prematurity, low birthweight and pre-eclampsia : an updated review with meta-analysis 2013;70:213–22. doi:10.1136/oemed-2012-101032.Work.
- [8] Zawilska JB, Skene DJ, Arendt J. Physiology and pharmacology of melatonin in relation to biological rhythms. *Pharmacol Reports* 2009;61:383–410. doi:10.1016/S1734-1140(09)70081-7.
- [9] Arendt J. Melatonin: Characteristics, Concerns, and Prospects. *J Biol Rhythms* 2005;20:291–303. doi:10.1177/0748730405277492.
- [10] Saarela S, Reiter RJ. Function of melatonin in thermoregulatory processes. *Life Sci* 1994;54:295–311. doi:10.1016/0024-3205(94)00786-1.
- [11] Maestroni GJM, Conti A. The pineal neurohormone melatonin stimulates activated CD4+, Thy-1+ cells to release opioid agonist(s) with immunoenhancing and anti-stress properties. *J Neuroimmunol* 1990;28:167–76. doi:10.1016/0165-5728(90)90031-H.
- [12] Poeggeler B, Thuermann S, Dose A, Schoenke M, Burkhardt S, Hardeland R. Melatonin's unique radical scavenging

- properties - Roles of its functional substituents as revealed by a comparison with its structural analogs. *J Pineal Res* 2002;33:20–30. doi:10.1034/j.1600-079X.2002.01873.x.
- [13] Hill SM, Teplitzky S, Ram PT, Kiefer T, Blask DE, Spriggs LL, et al. Melatonin synergizes with retinoic acid in the prevention and regression of breast cancer. *Adv Exp Med Biol* 1999;460:345–62.
- [14] Kiefer T, Ram PT, Yuan L, Hill SM. Melatonin inhibits estrogen receptor transactivation and cAMP levels in breast cancer cells. *Breast Cancer Res Treat* 2002;71:37–45. doi:10.1023/A:1013301408464.
- [15] Saper CB, Lu J, Chou TC, Gooley J. The hypothalamic integrator for circadian rhythms. *Trends Neurosci* 2005;28:152–7. doi:10.1016/j.tins.2004.12.009.
- [16] Dumont M, Lanctt V, Cadieux-Viau R, Paquet J. Melatonin production and light exposure of rotating night workers. *Chronobiol Int* 2012. doi:10.3109/07420528.2011.647177.
- [17] Guyon A, Balbo M, Morselli LL, Tasali E, Leproult R, L’Hermite-Balériaux M, et al. Adverse effects of two nights of sleep restriction on the hypothalamic-pituitary-adrenal axis in healthy men. *J Clin Endocrinol Metab* 2014;99:2861–8. doi:10.1210/jc.2013-4254.
- [18] Dugaard S, Garde AH, Bonde JPE, Christoffersen J, Hansen ÅM, Markvart J, et al. Night work, light exposure and melatonin on work days and days off. *Chronobiol Int* 2017. doi:10.1080/07420528.2017.1327867.
- [19] Rollag MD, Welsh MG, Klein DC. Effect of Acute Light Exposure Upon Melatonin Content, NAT Activity, and Nuclear Volume in the Gerbil Pineal Complex. *J Pineal Res* 1984;1:339–47. doi:10.1111/j.1600-079X.1984.tb00224.x.
- [20] Vartanian G V., Li BY, Chervenak AP, Walch OJ, Pack W, Ala-Laurila P, et al. Melatonin suppression by light in humans is more sensitive than previously reported. *J Biol Rhythms* 2015;30:351–4. doi:10.1177/0748730415585413.
- [21] Costa G, Akerstedt T, Nachreiner F, Balthieri F, Carvalhais J, Folkard S, et al. Flexible working hours, health, and well-being in Europe: Some considerations from a SALSTA project. *Chronobiol Int* 2004;21:831–44. doi:10.1081/LCBI-200035935.
- [22] Costa G, Di Milia L. Introductory overview - 19th international symposium on shiftwork and working time: Health and well-being in the 24-h society. *Chronobiol Int* 2010;27:889–97. doi:10.3109/07420528.2010.488606
- [23] Harrington JM. Health effects of shift work and extended hours of work. *Occup Environ Med* 2001. doi:10.1136/oem.58.1.68
- [24] Costa G. Sleep deprivation due to shift work. vol 131.1<sup>st</sup> ed.Elsevier B.V.;2015.doi:10.1016/B978-0-444-62627-1.00023-8
- [25] Erren TC, Reiter RJ, Piekarski C. Light, timing of biological rhythms, and chronodisruption in man. *Naturwissenschaften* 2003;90:485–94. doi:10.1007/s00114-003-0468-6.

- [26] Moreno CRDC, Fischer FM, Rotenberg L. A saúde do trabalhador na sociedade 24 horas. São Paulo Em Perspect 2003;17:34–46. doi:10.1590/S0102-88392003000100005.
- [27] Beersma DGM, Gordijn MCM. Circadian control of the sleep-wake cycle. *Physiol Behav* 2007;90:190–5. doi:10.1016/j.physbeh.2006.09.010.
- [28] Weaver DR. The Suprachiasmatic Nucleus: A 25-Year Retrospective. *J Biol Rhythms* 1998;13:100–12. doi:10.1177/074873098128999952.
- [29] Damiola F, Le Minli N, Preitner N, Kornmann B, Fleury-Olela F, Schibler U. Restricted feeding uncouples circadian oscillators in peripheral tissues from the central pacemaker in the suprachiasmatic nucleus. *Genes Dev* 2000;14:2950–61. doi:10.1101/gad.183500.
- [30] Morris CJ, Aeschbach D, Scheer FAJL. Circadian system, sleep and endocrinology. *Mol Cell Endocrinol* 2012. doi:10.1016/j.mce.2011.09.003.
- [31] Wehr TA, Aeschbach D, Jr WCD. Evidence for a biological dawn and dusk in the human circadian timing system. - PubMed - NCBI 2001:937–51.
- [32] Scheer FAJL, Hu K, Evoniuk H, Kelly EE, Malhotra A, Hilton MF, et al. Impact of the human circadian system, exercise, and their interaction on cardiovascular function. *Proc Natl Acad Sci* 2010;107:20541–6. doi:10.1073/pnas.1006749107.
- [33] Knutsson A, Boggild H. Shiftwork and cardiovascular disease: review of disease mechanisms. *Rev Environ Health* 2000;15:359–72.
- [34] Karlsson B, Knutsson A, Lindahl B. Is there an association between shift work and having a metabolic syndrome? Results from a population based study of 27,485 people. *Occup Environ Med* 2001;58:747–52. doi:10.1136/oem.58.11.747.
- [35] De Bacquer D, Van Risseghem M, Clays E, Kittel F, De Backer G, Braeckman L. Rotating shift work and the metabolic syndrome: a prospective study. *Int J Epidemiol* 2009;38:848–54. doi:10.1093/ije/dyn360.
- [36] Marqueze EC, Ulhôa MA, Moreno CRC. Irregular working times and metabolic disorders among truck drivers: A review. *Work* 2012;41:3718–25. doi:10.3233/WOR-2012-0085-3718.
- [37] Pietroiusti A, Neri A, Somma G, Coppeta L, Iavicoli I, Bergamaschi A, et al. Incidence of metabolic syndrome among night-shift healthcare workers. *Occup Environ Med* 2010;67:54–7. doi:10.1136/oem.2009.046797.
- [38] Biggi N, Consonni D, Galluzzo V, Sogliani M, Costa G. Metabolic syndrome in permanent night workers. *Chronobiol Int* 2008;25:443–54. doi:10.1080/07420520802114193.
- [39] Åkerstedt T. Shift work and disturbed sleep/wakefulness. *Occup Med (Chic Ill)* 2003;53:89–94.

doi:10.1093/ocmed/kg046.

- [40] Ketchum ES, Morton JM. Disappointing weight loss among shift workers after laparoscopic gastric bypass surgery. *Obes Surg* 2007;17:581–4. doi:10.1007/s11695-007-9100-8.
- [41] Morikawa Y, Nakagawa H, Miura K, Soyama Y, Ishizaki M, Kido T, et al. Effect of shift work on body mass index and metabolic parameters. *Scand J Work Environ Health* 2007;33:45–50. doi:10.5271/sjweh.1063.
- [42] Lowden A, Moreno C, Holmbäck U, Lennernäs M, Tucker P. Eating and shift work - Effects on habits, metabolism, and performance. *Scand J Work Environ Heal* 2010;36:150–62. doi:10.5271/sjweh.2898.
- [43] Bøggild H, Jeppesen HJ. Intervention in shift scheduling and changes in biomarkers of heart disease in hospital wards. *Scand J Work Environ Heal* 2001;27:87–96. doi:10.5271/sjweh.594.
- [44] Hakola T, Paukkonen M, Pohjonen T. Less quick returns--greater well-being. *Ind Health* 2010;48:390–4. doi:JST.JSTAGE/indhealth/MSSW-02 [pii].
- [45] Vyas M V., Garg AX, Iansavichus A V., Costella J, Donner A, Laugsand LE, et al. Shift work and vascular events: systematic review and meta-analysis. *Bmj* 2012;345:e4800–e4800. doi:10.1136/bmj.e4800.
- [46] Silva-Costa A, Rotenberg L, Nobre AA, Schmidt MI, Chor D, Griep RH. Gender-specific association between night-work exposure and type-2 diabetes: results from longitudinal study of adult health, ELSA-Brasil. *Scand J Work Environ Health* 2015;41:569–78. doi:10.5271/sjweh.3520.
- [47] Wang F, Zhang L, Zhang Y, Zhang B, He Y, Xie S, et al. Meta-analysis on night shift work and risk of metabolic syndrome. *Obes Rev* 2014;15:709–20. doi:10.1111/obr.12194.
- [48] Proper KI, Van De Langenberg D, Rodenburg W, Vermeulen RCH, Van Der Beek AJ, Van Steeg H, et al. The relationship between shift work and metabolic risk factors: A systematic review of longitudinal studies. *Am J Prev Med* 2016;50:e147–57. doi:10.1016/j.amepre.2015.11.013.
- [49] Schernhammer ES, Laden F, Speizer FE, Willett WC, Hunter DJ, Kawachi I, et al. Rotating Night Shifts and Risk of Breast Cancer in Women Participating in the Nurses' Health Study. *JNCI J Natl Cancer Inst* 2001;93:1563–8. doi:10.1093/jnci/93.20.1563.
- [50] Lee A, Myung S-K, Cho JJ, Jung Y-J, Yoon JL, Kim MY. Night Shift Work and Risk of Depression: Meta-analysis of Observational Studies. *J Korean Med Sci* 2017;32:1091–6. doi:10.3346/jkms.2017.32.7.1091.
- [51] Stocker LJ, MacKlon NS, Cheong YC, Bewley SJ. Influence of shift work on early reproductive outcomes: A systematic review and meta-analysis. *Obstet Gynecol* 2014;124:99–110. doi:10.1097/AOG.0000000000000321.

- [52] Haines VY, Marchand A, Rousseau V, Demers A. The mediating role of work-to-family conflict in the relationship between shiftwork and depression. *Work Stress* 2008;22:341–56. doi:10.1080/02678370802564272.
- [53] Richter HG, Torres-Farfrí C, Rojas-García PP, Campino C, Torrealba F, Serón-Ferré M. The circadian timing system: Making sense of day/night gene expression. *Biol Res* 2004;37:11–28. doi:10.4067/S0716-97602004000100003.
- [54] Amaral FG, Castrucci AM, Cipolla-Neto J, Poletini MO, Mendez N, Richter HG, et al. Environmental control of biological rhythms: Effects on development, fertility and metabolism. *J Neuroendocrinol* 2014;26:603–12. doi:10.1111/jne.12144.
- [55] Reiter RJ, Tan D-X, Manchester LC, Paredes SD, Mayo JC, Sainz RM. Melatonin and Reproduction Revisited. *Biol Reprod* 2009;81:445–56. doi:10.1095/biolreprod.108.075655.
- [56] Pévet P, Agez L, Bothorel B, Saboureaux M, Gauer F, Laurent V, et al. Melatonin in the multi-oscillatory mammalian circadian world. *Chronobiol Int* 2006;23:39–51. doi:10.1080/07420520500482074.
- [57] Johnston JD, Skene DJ. Regulation of mammalian neuroendocrine physiology and rhythms by melatonin. *J Endocrinol* 2015;226:T187–98. doi:10.1530/JOE-15-0119.
- [58] Cavallo A. Melatonin and human puberty: Current perspectives. *J Pineal Res* 1993;15:115-21. doi:10.1111/j.1600-079x.1993.tb00517.x.
- [59] Lampiao F. New developments of the effect of melatonin on reproduction. *World J Obstet Gynecol* 2013;2:15. doi:10.5317/wjog.v2.i2.15.
- [60] Silman RE. Melatonin : a contraceptive for the nineties. *Eur J Obstet Gynecol Reprod Biol* 1993;49:3-9. doi:10.1016/0028-2243(93)90099-x.
- [61] Tamura H, Nakamura Y, Terron MP, Flores LJ, Manchester LC, Tan DX, et al. Melatonin and pregnancy in the human. *Reprod Toxicol* 2008. doi:10.1016/j.reprotox.2008.03.005.
- [62] Landgraf D, Koch CE, Oster H. Embryonic development of circadian clocks in the mammalian suprachiasmatic nuclei. *Front Neuroanat* 2014. doi:10.3389/fnana.2014.00143.
- [63] Voiculescu SE, Zygouropoulos N, Zahi CD, Zagrean AM. Role of melatonin in embryo fetal development. *J Med Life* 2014;7:488–92. doi:25713608.
- [64] Ferreira DS, Amaral FG, Mesquita CC, Barbosa APL, Lellis-Santos C, Turati AO, et al. Maternal melatonin programs the daily pattern of energy metabolism in adult offspring. *PLoS One* 2012. doi:10.1371/journal.pone.0038795.
- [65] Knutsson A. Health disorders of shift workers. *Occup Med (Chic Ill)* 2003;53:103–8. doi:10.1093/occmed/kqg048.
- [66] Croteau A, Marcoux S, Brisson C. Work activity in pregnancy, preventive measures, and the risk of delivering a small-for-

- gestational-age infant. *Am J Public Health* 2006;96:846–55. doi:10.2105/AJPH.2004.058552.
- [67] Serón-Ferré M, Mendez N, Abarzua-Catalan L, Vilches N, Valenzuela FJ, Reynolds HE, et al. Circadian rhythms in the fetus. *Mol Cell Endocrinol* 2012;349:68–75. doi:10.1016/j.mce.2011.07.039.
- [68] Vilches N, Spichiger C, Mendez N, Abarzua-Catalan L, Galdames HA, Hazlerigg DG, et al. Gestational chronodisruption impairs hippocampal expression of NMDA receptor subunits Grin1b/Grin3a and spatial memory in the adult offspring. *PLoS One* 2014;9:17–9. doi:10.1371/journal.pone.0091313.
- [69] Motta-Teixeira LC, Machado-Nils AV, Battagello DS, Diniz GB, Andrade-Silva J, Silva S, et al. The absence of maternal pineal melatonin rhythm during pregnancy and lactation impairs offspring physical growth, neurodevelopment, and behavior. *Horm Behav* 2018;105:146–56. doi:10.1016/j.yhbeh.2018.08.006.
- [70] Gluckman PD. Effect of In Utero and Early Life Conditions on Adult Health and Disease. *N Engl J Med* 2008;359:61–73. doi:10.1056/NEJMra0708473.Effect.
- [71] Reiter RJ. Melatonin and human reproduction. *Ann Med* 1998;30:103–8. doi:10.3109/07853899808999391.
- [72] Serón-Ferré M, Ducsay CA, Valenzuela GJ. Circadian rhythms during pregnancy. *Endocr Rev* 1993;14:594–609. doi:10.1210/edrv-14-5-594.
- [73] Deguchi T. Ontogenesis of a biological clock for serotonin:acetyl coenzyme A N-acetyltransferase in pineal gland of rat (circadian/neural regulation/melatonin). *Physiology* 1975;72:2814–8.
- [74] Reppert SM, Chez RA, Anderson A, Klein DC. Maternal-fetal transfer of melatonin in the non-human primate. *Pediatr Res* 1979;13:788–91. doi:10.1203/00006450-197906000-00015.
- [75] Yellon SM, Longo LD. Melatonin rhythms in fetal and maternal circulation during pregnancy in sheep. *Am J Physiol* 1987;252:E799-802. doi:10.1152/ajpendo.1987.252.6.E799.
- [76] Serón-Ferré M, Torres-Farfán C, Forcelledo ML, Valenzuela GJ. The development of circadian rhythms in the fetus and neonate. *Semin Perinatol* 2001;25:363–70. doi:10.1053/sper.2001.29037.
- [77] Mirmiran M, Maas YGH, Ariagno RL. Development of fetal and neonatal sleep and circadian rhythms. *Sleep Med Rev* 2003;7:321–34. doi:10.1053/smr.2002.0243.
- [78] Patrick J, Campbell K, Carmichael L, Natale R, Richardson B. Patterns of gross fetal body movements over 24-hour observation intervals during the last 10 weeks of pregnancy. *Am J Obstet Gynecol* 1982;142:363–71. doi:10.1016/S0002-9378(16)32375-4.
- [79] Cipolla-Neto J, Amaral FG, Afeche SC, Tan DX, Reiter RJ. Melatonin, energy metabolism, and obesity: A review. vol. 56.

2014. doi:10.1111/jpi.12137.
- [80] Reiter RJ, Tan DX, Korkmaz A, Rosales-Corral SA. Melatonin and stable circadian rhythms optimize maternal, placental and fetal physiology. *Hum Reprod Update* 2014;20:293–307. doi:10.1093/humupd/dmt054.
- [81] Lucas A, Fewtrell MS, Cole TJ. Fetal origins of adult disease—the hypothesis revisited. *Bmj* 1999;319:245. doi:10.1136/bmj.319.7204.245.
- [82] Iliadou A. Low birthweight and Type 2 diabetes: A study on 11 162 Swedish twins. *Int J Epidemiol* 2004;33:948–53. doi:10.1093/ije/dyh117.
- [83] Whitaker R, Wright J, Pepe M, Seidel K, Dietz W. Predicting Obesity in Young Adulthood From Childhood. *Massachusetts Med Soc* 1997;337:869–73. doi:10.1056/NEJM199709253371301.
- [84] Ross MG, Desai M. Developmental programming of offspring obesity, adipogenesis, and appetite. *Clin Obstet Gynecol* 2013;56:529–36. doi:10.1097/GRF.0b013e318299c39d.
- [85] Diaz B, Blazquez E. Effect of Pinealectomy on Plasma Glucose, Insulin and Glucagon Levels in the Rat. *Horm Metab* 1986;18:225–9.
- [86] Lima FB, Machado UF, Bartol I, Seraphim PM, Sumida DH, Moraes SM, et al. Pinealectomy causes glucose intolerance and decreases adipose cell responsiveness to insulin in rats. *Am J Physiol* 1998;275:E934–41.
- [87] Anhê GF, Caperuto LC, Pereira-Da-Silva M, Souza LC, Hirata AE, Velloso LA, et al. In vivo activation of insulin receptor tyrosine kinase by melatonin in the rat hypothalamus. *J Neurochem* 2004. doi:10.1111/j.1471-4159.2004.02514.x.
- [88] Picinato MC, Hirata AE, Cipolla-Neto J, Curi R, Carvalho CRO, Anhê GF, et al. Activation of insulin and IGF-1 signaling pathways by melatonin through MT1 receptor in isolated rat pancreatic islets. *J Pineal Res* 2008;44:88–94. doi:10.1111/j.1600-079X.2007.00493.x.
- [89] Van Cauter E, Blackman JD, Roland D, Spire JP, Refetoff S, Polonsky KS. Modulation of glucose regulation and insulin secretion by circadian rhythmicity and sleep. *J Clin Invest* 1991;88:934–42. doi:10.1172/JCI115396.
- [90] Van Cauter E, Polonsky KS, Scheen AJ. Roles of circadian rhythmicity and sleep in human glucose regulation. *Endocr Rev* 1997;18:716–38. doi:10.1210/er.18.5.716.
- [91] Scheer FAJL, Hilton MF, Mantzoros CS, Shea SA. Adverse metabolic and cardiovascular consequences of circadian misalignment. *Proc Natl Acad Sci* 2009;106:4453–8. doi:10.1073/pnas.0808180106.
- [92] Bizot-Espiard JG, Double A, Guardiola-Lemaitre B, Delagrangre P, Ktorza A, Penicaud L. Diurnal rhythms in plasma glucose, insulin, growth hormone and melatonin levels in fasted and hyperglycaemic rats. *Diabetes Metab* 1998.



- [93] la Fleur SE, Kalsbeek a, Wortel J, van der Vliet J, Buijs RM. Role for the pineal and melatonin in glucose homeostasis: pinealectomy increases night-time glucose concentrations. *J Neuroendocrinol* 2001. doi:10.1046/j.1365-2826.2001.00717.x.
- [94] Gottlieb DJ, Punjabi NM, Newman AB, Resnick HE, Redline S, Baldwin CM, et al. Association of sleep time with diabetes mellitus and impaired glucose tolerance. *Arch Intern Med* 2005. doi:10.1001/archinte.165.8.863.
- [95] Donga E, van Dijk M, van Dijk JG, Biermasz NR, Lammers G-J, van Kralingen KW, et al. A single night of partial sleep deprivation induces insulin resistance in multiple metabolic pathways in healthy subjects. *J Clin Endocrinol Metab* 2010. doi:10.1210/jc.2009-2430.
- [96] Ramracheya RD, Muller DS, Squires PE, Brereton H, Sugden D, Huang GC, et al. Function and expression of melatonin receptors on human pancreatic islets. *J Pineal Res* 2008;44:273–9. doi:10.1111/j.1600-079X.2007.00523.x.
- [97] Masana MI, Dubocovich ML. Melatonin Receptor Signaling: Finding the Path Through the Dark. *Sci STKE* 2001;107:39–39. doi:10.1126/stke.2001.107.pe39.
- [98] Alonso-Vale MIC, Andreotti S, Peres SB, Anhê GF, das Neves Borges-Silva C, Neto JC, et al. Melatonin enhances leptin expression by rat adipocytes in the presence of insulin. *Am J Physiol Metab* 2005;288:E805–12. doi:10.1152/ajpendo.00478.2004.
- [99] Gibson T, Stimmler L, Jarrett RJ, Rutland P, Shiu M. Diurnal variation in the effects of insulin on blood glucose, plasma non-esterified fatty acids and growth hormone. *Diabetologia* 1975;11:83–8. doi:10.1007/BF00422823.
- [100] Montano ME, Molpeceres V, Mauriz JL, Garzo E, Cruz IBM, González P, et al. Effect of melatonin supplementation on food and water intake in streptozotocin-diabetic and non-diabetic male Wistar rats. *Nutr Hosp* 2010;25:931–8. doi:10.3305/nh.2010.25.6.4803.
- [101] Zanuto R, Siqueira-Filho MA, Caperuto LC, Bacurau RFP, Hirata E, Peliciari-Garcia RA, et al. Melatonin improves insulin sensitivity independently of weight loss in old obese rats. *J Pineal Res* 2013;55:156–65. doi:10.1111/jpi.12056.
- [102] Barrios Y, Díaz N, Meertens L, Naddaf G, Solano L, Fernández M, et al. Leptina sérica, su relación con peso y distribución de grasa corporal en mujeres posmenopáusicas. *Nutr Hosp* 2010;25:80–4. doi:10.3305/nh.2010.25.1.4273.
- [103] Kojima S, Nakahara T, Nagai N, Muranaga T, Tanaka M, Yasuhara D, et al. Altered ghrelin and peptide YY responses to meals in bulimia nervosa. *Clin Endocrinol (Oxf)* 2005;62:74–8. doi:10.1111/j.1365-2265.2004.02176.x.
- [104] Cambuli VM, Musiu MC, Incani M, Paderi M, Serpe R, Marras V, et al. Assessment of adiponectin and leptin as biomarkers of positive metabolic outcomes after lifestyle intervention in overweight and obese children. *J Clin Endocrinol Metab* 2008;93:3051–7. doi:10.1210/jc.2008-0476.

- [105] Wehr TA. Photoperiodism in humans and other primates: Evidence and implications. *J Biol Rhythms* 2001;16:348–64.  
doi:10.1177/074873001129002060.
- [106] Haus E, Smolensky M. Biological clocks and shift work: Circadian dysregulation and potential long-term effects. *Cancer Causes Control* 2006; 17:489-500.doi:10.1007/s10552-005-9015-4.